

- the T-cell receptor prevents autoimmune encephalomyelitis and activates Th2 immunity. *Nature Medicine* 2:899-905.
42. Ramierz, F., D. J. Fowell, M. Puklavec, S. Simmonds, and D. Mason. 1996. Glucocorticoids promote a TH2 cytokine response by CD4+ T cells in vitro. *J Immunol* 156:2406-12.
  43. Saoudi, A., S. Simmonds, I. Huitinga, and D. Mason. 1995. Prevention of experimental allergic encephalomyelitis in rats by targeting autoantigen to B cells: evidence that the protective mechanism depends on changes in the cytokine response and migratory properties of the autoantigen-specific T cells [see comments]. *J Exp Med* 182:335-44.
  44. Thornhill, M. H., S. M. Wellicome, D. L. Mahiouz, J. S. Lanchbury, A. U. Kyan, and D. O. Haskard. 1991. Tumor necrosis factor combines with IL-4 or IFN-gamma to selectively enhance endothelial cell adhesiveness for T cells. The contribution of vascular cell adhesion molecule-1-dependent and -independent binding mechanisms. *J Immunol* 146:592-8.
  45. Baron, J. L., J. A. Madri, N. H. Ruddle, G. Hashim, and C. J. Janeway. 1993. Surface expression of alpha 4 integrin by CD4 T cells is required for their entry into brain parenchyma. *J Exp Med* 177:57-68.
  46. Yednock, T. A., C. Cannon, L. C. Fritz, M. F. Sanchez, L. Steinman, and N. Karin. 1992. Prevention of experimental autoimmune encephalomyelitis by antibodies against alpha 4 beta 1 integrin. *Nature* 356:63-6.
  47. Kuroda, Y., and Y. Shimamoto. 1991. Human tumor necrosis factor-alpha augments experimental allergic encephalomyelitis in rats. *J Neuroimmunol* 34:159-64.
  48. Powell, M. B., D. Mitchell, J. Lederman, J. Buckmeier, S. S. Zamvil, M. Graham, N. H. Ruddle, and L. Steinman. 1990. Lymphotoxin and tumor necrosis factor-alpha production by myelin basic protein-specific T cell clones correlates with encephalitogenicity. *Int Immunol* 2:539-44.
  49. Ruddle, N. H., C. M. Bergman, K. M. McGrath, E. G. Lingenheld, M. L. Grunnet, S. J. Padula, and R. B. Clark. 1990. An antibody to lymphotoxin and tumor necrosis factor prevents transfer of experimental allergic encephalomyelitis. *J Exp Med* 172:1193-200.
  50. Selmaj, K., C. S. Raine, and A. H. Cross. 1991. Anti-tumor necrosis factor therapy abrogates autoimmune demyelination. *Ann Neurol* 30:694-700.

51. Kohno, K., J. Kataoka, T. Ohtsuki, Y. Suemoto, I. Okamoto, M. Usui, M. Ikeda, and M. Kurimoto. 1997. IFN-gamma-inducing factor (IGIF) is a costimulatory factor on the activation of Th1 but not Th2 cells and exerts its effect independently of IL-12. *J Immunol* 158:1541-50.
52. Rothe, H., N. A. Jenkins, N. G. Copeland, and H. Kolb. 1997. Active stage of autoimmune diabetes is associated with the expression of a novel cytokine, IGIF, which is located near Idd2. *J Clin Invest* 99:469-74.
53. Karin, N., J. D. Mitchell, S. Brocke, N. Ling, and L. Steinman. 1994. Reversal of experimental autoimmune encephalomyelitis by a soluble peptide variant of a myelin basic protein epitope: T cell receptor antagonism and reduction of IFN-g and TNF-a production. *J. Exp. Med.* 180:2227-2237.
54. Grewal, I. S., H. G. Foellmer, K. D. Grewal, J. Xu, F. Hardardottir, J. L. Baron, C. A. Janeway, Jr., and R. A. Flavell. 1996. Requirement for CD40 ligand in costimulation induction, T cell activation, and experimental allergic encephalomyelitis. *Science* 273:1864-7.
55. Krakowski, M., and T. Owens. 1996. Interferon-gamma confers resistance to experimental allergic encephalomyelitis. *Eur J Immunol* 26:1641-6.
56. Heremans, H., C. Dillen, M. Groenen, E. Martens, and A. Billiau. 1996. Chronic relapsing experimental autoimmune encephalomyelitis (CREAE) in mice: enhancement by monoclonal antibodies against interferon-gamma. *Eur J Immunol* 26:2393-8.
57. Billiau, A., H. Heremans, F. Vandekerckhove, R. Dijkmans, H. Sobis, E. Meulepas, and H. Carton. 1988. Enhancement of experimental allergic encephalomyelitis in mice by antibodies against IFN-gamma. *J Immunol* 140:1506-10.
58. Segal, B. M., B. K. Dwyer, and E. M. Shevach. 1998. An interleukin (IL)-10/IL-12 immunoregulatory circuit controls susceptibility to autoimmune disease. *J Exp Med* 187:537-46.

1. An antibody comprising an immunoglobulin capable of binding an interferon gamma inducing factor.
2. The antibody of claim 1, wherein the antibody is polyclonal.
3. The antibody of claim 1, wherein the antibody is monoclonal.
4. The antibody of claim 1, wherein said antibody is a neutralizing antibodies to said interferon gamma inducing factor in affecting cells to produce interferon gamma.
5. The antibody of claim 1, wherein the antibody is humanized.
6. A method for treating an animal for inducing protective immunity against multiple sclerosis, the method comprising the step of administering to said animal cells being capable of producing and secreting an antibody capable of neutralizing an interferon gamma inducing factor in affecting cells to produce interferon gamma.
7. The method of claim 6, wherein the antibody is humanized.
8. A method for treating an animal for inducing protective immunity against multiple sclerosis, the method comprising the step of administering to said animal an antibody capable of neutralizing an interferon gamma inducing factor in affecting cells to produce interferon gamma.
9. The method of claim 8, wherein the antibody is humanized.
10. A pharmaceutical composition for inducing protective immunity against multiple sclerosis, comprising a pharmaceutically acceptable carrier and an antibody being capable of binding an interferon gamma inducing factor.

09856798-08401  
T04280-8629880